



Birds on Top of 4/6 83.

*Larus cristatus*

*Grus columba*  
*Phalaropus lobatus*  
*Phalaropus lobatus*

*Phalaropus cristatus*

*Grus cristatus*

*Fulica marila*

~~*Grus cristatus*~~ - 5 eggs?

*Grus cristatus*

*Grus cristatus*

*Grus cristatus* 3 eggs.

*Grus cristatus*

*Grus cristatus*

*Grus cristatus*

Birds at other locations


*Grus cristatus*

*Grus cristatus*

*Grus cristatus*

*Grus cristatus*

Dimensions of site for Top of 4.9-1.0 m.  
Length of site 4.0 m.



Storoccephalus dubovskii

(no 2184)  
Oscha shot one (2/683 between Luvina  
and Saravuma; it was in company with  
two of the same kind.

Chelodactylus borealis

1883 No. 1000 at the village 8/10. - The  
following day an other. - 1 1/2 two from  
Luvina, where they were said to be  
common. (The same) 3 shot at 17. -  
1 1/2 shot at 17, and one shot at 17, same  
day. The same as from Luvina. 2 1/2  
and one from D.D. -

Turdus obscurus?

One ♂, which Osler shot 10/6/83 was  
wounded about a week ago. before that  
date. - Another was shot 11/6/83. - 12/6/83.  
13<sup>th</sup> was shot at Plover. 15<sup>th</sup> was shot at  
Govan. \* and one at Gavan. - 7<sup>th</sup> was shot at  
Govan.

Merula pallascus Linn.

One ♂ shot.  
One ♂ shot at Gavan. 10/6/83.  
One ♂ shot at Gavan. 11/6/83. - 12/6/83.  
One ♂ shot at Gavan. 15/6/83. - 16/6/83.



March 1894  
Sun 7 11.

Chas. H. H. H.

Actacilla melanogaster

Found - 3/5 - near Tsim 4/580 and  
also the following day, when I got two -  
11/5 saw two individuals. - 12/5 saw two  
about 1300 ft of June

They were seen near the landing this year  
and as it was kept low into the water  
of the island. 4/580 about 10 miles S.W. of A.D.

*From Refugio*

Actacilla haemorrhagica

The first was 1883 at Ref 6/583 - to 20108. This  
specimen was seen about 4/5. - 5/583. at 1000 ft  
at Tsim 6. D. 2 - ~~from Tsim~~

None have been seen <sup>(10/583)</sup> since my return  
from Otago 31/583 -

*Podiceps flavus*  
Saw many very abundant in Saginaw Lake  
about water shot on May 21, 22<sup>nd</sup>, 23<sup>rd</sup>, 24<sup>th</sup>  
and 31<sup>st</sup> of May 1883. Another was shot at  
Ladysmith 3<sup>rd</sup> June 83. - Observed two  
in Garfield. Tuesday 5/6. - Two seen  
and one shot 10/6 83. -

*Podiceps flavus*



Chelidon Hepuensis Ridgw.

3/6 83 one seen, but has probably arrived long ago, as the eggs obtained 3/11 83 probably belong to that species. - Met fully fledged young at Karabuluj, Ceylon Is., 9/7 83, and got a nest with two eggs at Glinka 13/7 83, with two seen to breed twice. -

Besides the <sup>common</sup> chirping and twitter when it has a short song tsip, tsip, tsip, and its call note is a short ~~tsip~~ tsip tsip!

Chelidon Hepuensis  
i 15/9 83. 2 seen at Karabuluj, Ceylon Is. -  
also seen at Glinka Is. -

Erythronema leucura?

As I in the morning 83 got  
the first specimen on board the steamer  
Alexander when on the way from Petro-  
paulsk to Derling Island, and showed  
it to Dr. Dybowski, he asked me, whether  
this species ~~did~~ occurs on the island  
or not. As I had never seen or got  
it there, I answered, that it did  
not occur, having no idea of - of  
course - that a ♀ was shot on  
the island in the very hour in which  
this conversation took place.

Muscicapa  
tricolor

A large number must have arrived  
yesterday or last night (8/683) as about  
10-12 were observed, and 6 shot. Yester-  
day the wind was S and changed during  
the night to E with variable force from  
very light to fresh. - On following day  
we saw at least 9 and shot 5. - The 10<sup>th</sup>  
I were shot, and several seen. - However  
on all places sheltered from the high fresh  
wind of today (11/683), I myself caught  
more than 20, from near shot. - The birds  
seen. - In the last two days birds have not  
been scarce here at Gorka, but today (12/6)  
Kargan brought some from Srednaya that he  
a shot on the way. - The same day I shot  
shot at Gorka and brought away the 10<sup>th</sup> at  
19/683. - I also brought one shot yesterday from  
the same place. Several seen. -

Chelidon saturata

Cocher states, that he saw two swallows, black above and reddish brown below at the North Cookey D. I. 19/6 83, but as they were too near the seals it was impossible to secure them. - 22/6 83 old Nurdukowski saw one swallow, black above and below, at Ladiginsk. -

Fringilla montifringilla

Rather numerous this ~~day~~ spring; besides the 5 ones prepared, 3 others were received about 1/6 83. -

*Linaria fucata*

*Alcedo gularis*

Exp. 26, Karatschi 7/783 shot three.  
fully fledged young in one shot. It  
stands as if the parent were for a  
second time, as I have met several  
pairs in that position of their legs. —  
They are usually found only along the steep  
rocks of the ~~the~~ coast, but sometimes also  
in the interior. Thus I met a family 10/783  
at a small creek in the interior of Copper  
Island at Karatschi; the note of the young  
is deeper and lower than that of the adults  
and is almost thrushlike. —



L. costata brunneica

1875-2



Acathis

Leptocarpus variabilis?

Shot one example ~~just before~~ in the narrow valley just behind. When the water supply of the village is. It was terribly injured by the shot (2 grain powder and shot No. 2), so that it represented only a large, single hole. One of both mandibles broken and the skull completely broken to pieces etc. It is a luck, that the color is so conspicuous, because the skin is only so good - or so bad - as the circumstances would allow -

— "Leptocarpus variabilis" Jacq. J. p. Am. 1881 p. 184  
from St. Paul and Kamtschatka. —  
Leptocarpus variabilis Kohl.

Hypocentrus aureolus

Leptocarpus variabilis

Phaethon rubricauda

1883. 13/3 83. Still only single birds. Commences  
to sing. - 18/3 birds heard along north beach. -  
28/3 the first flock consisting of 6 individuals  
at Reef. - 19/4 first set commenced to lay eggs  
at Reef. - Upper Is. natives assert, that  
they lay eggs twice in summer. -

Callaeus leucurus

1883. The first one taken on 21<sup>st</sup> of April; the first one shot the following day.

— 14<sup>th</sup>. A rather slender young bird. The bill was slender, and the throat white, and the wings of a light yellowish.

*Callaeus leucurus*

Alaska

19485 shot No. 10000 the last of the  
beach.





Ceculus canonicus

Brachyotus acipitrinus

4/583 Saw one at the old graveyard, N. I.

8/687 Looked through of this one without finding it.

*Brachyotus acipitrinus*

Isks met a family of 5 indiv. on the road  
between Severn Roakery and Gavan B. I. 3/983.

He says, that a pair was seen the whole summer  
in the neighborhood of the rookery, and that  
Grebittsky has got nest-youngs from Kamenev,  
some years ago. —

Microfates cardigans

How all from



Falls

Several young ones <sup>at a "Hot Springs"</sup> near Gavan, Copper Id.

Thalassidroma pacifica

First seen on Laysan about 1893. It was reported from Hawaii in the spring of 1894.

Halimastur

Old eggs

Lagopus

of 13 ind. shot 29/2 87, but had one or two dark feathers, as usual the trace of new feathers yet. - They are much more deficient to meet now than before, and while a month ago 30-40 were the average amount secured of half a day's shooting, it is now seldom that one obtains 10. - 17/4 88 shot 14 in a flight of 915 ft. the big flock out with some brown and feathers.

April - 6/7 83 found a nest close by the Blackburne house, at "Midjane looking; "Hale", "Buckley, etc." containing 9 eggs. Nest had a diameter of 140 mm consisting of dry grass and a few feathers. The eggs were small fresh.

Samaloua

A. Samaloua was shown to the study  
block at Lagoon 1983, but as they were very  
dry I could not reach any of them. The site  
did not seem flat but the black line from the  
ridge with a large white patch. This is  
probably a small piece of water. The ground  
is 2 ft. deep. The ground is very  
fine all over.

*Adipos testis*

Morinella integras

7/10/83 is said that Halimulus has got out today.

Squataroda histrica



Charadrius fuscus

1883. The first one was shot the 15<sup>th</sup> of May.  
June 2<sup>d</sup>. I observed half a dozen on the  
tundra and shot one. -

Chrysotis

Agialites alexandrinus

9883. Cyprus Is. Observed at least half a dozen  
on the about 1000' high ridge between Perithous and  
Kastellorizo.

*Agialites alexandrinus*

Eudynamis mongolicus

4/6 83. Found of 2. Shot with 3 eggs, 2 shot on the nest (no 2123) The eggs were lying with the points upwards and downwards, ~~as~~ in a slight hollow, ~~between~~ between four Agrostis arvensis, ~~with the~~ dry particles of leaves and other of the plant and numerous seeds of the same formed the nest; they were evidently brought together by the bird. Situation about 40 feet from ~~the~~ <sup>the</sup> sea and about 14 feet above the level of the sea. -- \* highest mark. One of the eggs was open. The bird contained an embryo with large eyes, the head only a little more developed than, with wings, legs etc.

4/10. 2 eggs. Found of 2. Shot with 3 eggs, 2 shot on the nest (no 2123) The eggs were lying with the points upwards and downwards, ~~as~~ in a slight hollow, ~~between~~ between four Agrostis arvensis, ~~with the~~ dry particles of leaves and other of the plant and numerous seeds of the same formed the nest; they were evidently brought together by the bird. Situation about 40 feet from ~~the~~ <sup>the</sup> sea and about 14 feet above the level of the sea. -- \* highest mark. One of the eggs was open. The bird contained an embryo with large eyes, the head only a little more developed than, with wings, legs etc.

12/7 83 at 10.30 am. Glimpses I was brought a not yet fully feathered young albatross. Allowed to run on the floor, it immediately commenced a very hot pursuit of the flies, which were caught with a remarkable precision and rapidity. It did not take any notice of the presence of man, but when the dog in the corner roared, it suddenly dropped flat on the floor with withdrawn neck, making itself as low as possible and rested so absolutely immovable until the dog had turned its back to it, when it ran off to any concealed place to hid itself. Within <sup>half</sup> an hour it had, however, made the experience that the dog did not ~~take any notice of~~ <sup>it</sup> ~~it~~, and consequently <sup>it</sup> paid no

little attention to the dog as to the man. (ms 2242).

Gallinago

1933<sup>33</sup> Vol. 1, No. 1, 1933, and still more —

Argemone

At home he will write in 1830-31.  
1833 shows a picture of a plant of not less  
than 50 inches. - being a small plant with  
spikes and small leaves. - the seed is small and  
is not so large as the seed of the last.

1833. It is not so large as the seed of the last.  
The seed is small and is not so large as the seed of the last.



Atakronus tepanickii

*Chrysophis flavus*

Setonius apicatus

Sceloporus sublineatus

Two shot of Gavanke Osw D-I. 27/6 83.  
brother of the same 22/6 83.

*Sceloporus sublineatus*

Helidius alpina

*Callis nivalis*

*Callis nivalis*



Muscicivora

One shot 20/6 and one 21/6 83 at (Severna), Emilia,  
Novski Mts, D. I. Was alone. —

*2. on 21/6 83, Severna, Emilia, Novski Mts, D. I.*

Osche met a flock on the plateau between  
Garan and Severna D. I. 3/983, feeding in the berries  
of Empetrum.

Lincoln's Sparrow-Clayton

one shot of which at not far from, at Fiddler's  
10/5/83, in gray plumage.

1 shot and prepared during my absence

One of two shot at Gavan N.P. 1/6/83, in red  
plumage.

Two shot at Severine 18/6/83.

One at Glinka, Copper Lk mid 7/83 and another  
at Pestchaniy, 30/7/83.

Totanus ochroleucus

Phaeocephalus glaucus

1 shot at Kaniuni 7/6 83.—

*Phaeocephalus glaucus*

Heteroscelus micans

Ocho saw and shot the first one B. 9. 83  $\frac{25}{5}$ ,  
during my absence in Otmuchowski.

In Cape Bar I saw one on the beach at Glukha  
14/4 83. - which was seen in the same place in 1882.  
1881 at the same place.



Aetideus hypoleucus

*Stellina*

*Limnodynastes acuminatus.*

*Limnodynastes* 1

Lobipes lobatus

During my absence in Peloponnesos  
observed by the first house at Demos, 2nd  
May 24<sup>th</sup> 1883. To 1<sup>st</sup> of June I found them  
numerous on the Thardos.

*Helicoverpa armigera*

Ola eggules

Found at ...  
...  
A ...  
... 24/4 84.

Chen — ?

Several white geese with black tipped  
wings are reported to have been seen on  
the surrounding lakes, northern part of Bering Id.  
5-6 Sept 1883. Said to be very shy.

Anser albifrons

*Anser albifrons*



Epanta leucoparva  
27-83 Oct 10 I & Synopsea Reef B-3  
Cruising from the sea.

Franta nigricans

*Helianthus laevis*

*Spizella clypeata*

A single pair, male observed between Colarus  
and Alaska. A 2. 24<sup>th</sup> of April 1883. Probably  
a few more birds in the Stathsonian part.  
Truslow's Konikowski reports, that they  
breed. One ♂ observed obtained 15 83<sup>4</sup>  
and one observed 5/6 83 on the Tundra Garment.

*Spizella clypeata*

Along the northern shore, between Fairbanks and  
Seward. S. A. and Seward. 1883.  
12/18/83 near Seward 3 eggs at Fairbanks  
19/83 from Seward 7 fresh eggs  
in Fairbanks.

Oafela acuta

23/4 83, a few ones observed between  
Thomson and Calverton.

27/4 83. Poole saw a small flock at Reef  
to day. 28/4 O. shot one at Loughwick.

4/6 83 Nest with 7 eggs on Top of Is.

10/6 83 a nest with 5 fresh eggs from Foully.

Spencer

Marsca penelope

Order Nov 4 & Bristol's St. Catherine  
to Nov 30/483. — ~~See then~~ The bees  
rather numerous this spring, and several  
were still seen 7/683.

*Cyathus penelope* *adignus*.



Station 1000

April 13<sup>th</sup> 1885, the first, a single one, was  
seen in a small flock of black-backed geese.  
Weather changed the morning from cold to mild  
with an E. wind. - Saw 2 or 3 others  
near the C. R. R. bridge to Lake Superior.

22-24<sup>th</sup> I saw common black birds along  
and around the lake.

Yokel was out of a Michigan

Yokel was out of a Michigan  
Yokel was out of a Michigan  
Yokel was out of a Michigan

*Quercus macrocarpa*

Ensetia falcata

Syn: "Ensis falcaria" Full. Amer. H. III p. 701 n. 34.

Quell. bot. N. I. II p. 521. n. 86

Ensis falcata Georgi Meisn. p. 167. - Pall.

Europ. II p. 259 n. 343 " ex Meisn.

Suppl. ad Europ. Ross. p. 21.

Ensetia falcata  
D. J. 1833. 33.

Fuligula cristata

7/683 a ♂ and a ♀ were shot in  
Kamensk, only the ♂ being fit for pres-  
ervation. Ochet thinks that he shot a ♂  
during my absence in Petropaulsk  
mistaking it for F. marila.

Fulix marila

5/583 the first at Gavran D. O. - by the  
first shot, a ♀.

*Fuligula cristata*

Arthropia

Mr. Osche describes a duck which  
he met 5/6 83 in Hannover, in compar-  
ison with Fuligula marila, as being  
exactly like the male of this species,  
but having a chestnut head. —

Clangula glaucion

Mr. Osche describes 32 1945 3 of 2 5

*Clangula glaucion*

Histrionetia minutus

1882. Aug. 10. at camp on the river.

11/283 a large flock at Ritz.

21/683 I saw met a large flock<sup>♂+♀</sup> in the  
small Creek Nolsjaja Rakuselnik, Severn.

Thylacantha Brandt (see Thylacantha)

Mem. Acad. Imp. Sc. St. Peterb. Vols. Sc. nat.

tom. six. (1848) - p. 4. Descrip.

φ η υ α ε & histrio & ν η τ τ α anas.

Chondrus fulvus



Harlequin hypomelae  
No 7 April 1882 in full change of plumage.

*Colymbus cristatus*

Enicometta stelleri

1883. 12/3 Large flocks of ♂♂ and ♀♀ at Frodhu  
1/4 83 numerous ♂♂ at do. - do. - do.  
2-5/83. 20-25/83 numerous flocks between  
Kauai & Tolu, N. P. - 28/4 83. Some  
few individuals at Reef, O. I. - Also a few 5/5 83 -  
4/5 83 A few. - 5/5 83 several ♂♂ and ♀♀ -  
5/5 83 half a dozen at Tolu, O. I. - 19/5 Large  
flocks between Kauai & Tolu, N. P. - Disappeared  
about 1/5 of May 83. -

*Enicometta stelleri*

Smilax virginica

# 83 a P. 1111 at bottom of Lycopodium B. 2

Smilax

*Melanetta americana*  
~~velutosa~~  
Makiston havde den fra Kautschatka.

*Melanetta fusca*

*Melanetta fusca*  
S. a. D. Inf. 9. 1. 1843.

*Mergus serrator*

Copper Pt. Petichanij Resolika, Gauran, 4883 a g.

*Mergus morganensis*  
P. 15/4883! L. by "Knechtli" was seen to day  
at Gauran 4883!

*Cyprinus carpio*



Larus glaucescens  
had eggs on the Lagoon 16/1-83. —  
Found a bird was brought from  
the same place 17/1-83. and of the same  
descent group.

Alca Laysana  
April 1st 1880. — the first flocks seen at  
Foster's Bay.  
First Young out 4883, Egg No. —

*Alca Laysana*

Chisa brevicauda  
April 7<sup>th</sup> 1883. - The first birds seen at  
Tolmie Falls.

*Chloris phoenicea*

*Stroma parvula* Mun.??  
the first ones seen and two others  
5/6 83 at the tundra at Gavan. 9/6 Osher saw  
4 there.

*Stroma longipennis*  
Besides the one shot several  
others were seen. 24/6 83 one got from Emilia,  
near the ship D. I.

29/6 83 plenty of them at Rakovaya  
Anchorage, Anzheba Bay, Kamt.  
Ch. Kottwitz ± p 322 *Stroma glacialis*,  
Lill = *S. kamtschatica* Penn & Fall.

*Stroma longipennis*

Sticovarus lanceolatus

4/6 83. One or two were observed  
when on Foxcroft Island. Got one  
the next day. - Also one 2/6 83 from  
the small lake at Marochinsk between  
Sarama and Gavan B.I. - One from  
Gladkovskij, Copper Is. 12 83, no  
~~then~~ saw another at Karabelnij same day after --  
At my arrival at Glinka I found there two  
specimens: tarsus blue, feet black. - In note  
at Gavan Sept 83 22/21.

Sticovarus parvulus

4/5 83 The first one at Bay A.I.  
1/6 83 An example of the light phase. 20/6 86.

Sticovarus lanceolatus

Diomedea bartholemica

1980 found the remains of an adult near  
Cape St. Lawrence. No more seen since that  
time.

Dr. Cook saw one at Cape 11/3/82, and not  
less than 8 white ones 11/4/82. —  
Saw one adult 2/5/83 at Cape 11.5.

Oceanodroma furcata

*Oceanodroma furcata*

Fulmarus

April 15 1981. On island around to my house. The  
large quantity of food like fish bones of *Stenobothrus*.



*Festuca ovina*

Emerson's Parody of Job was, ~~it~~ (1833) but  
he was 10 days ago had some of the same  
Carnegie A. I. - 1833 and was a lot more —

76 22. *Thela* brought one egg from Kamey;  
only one being in the nest. —

— 20/7 83 Cyper H. one pair at Portchara Oxen, Gauru.

Fabesula consociata

*Hypocentor auriculatus*

Lanius borealis

7/5/83. Starvo returned from an expedition to  
 the interior, during which he had observed  
 several specimens of this and several other  
 birds. The young is readily distinguished from the  
 adult by its smaller size, more brown and yellow  
 on the underparts, more dark and barred  
 with brown on the back, and a light  
 blue on the crown, more curved than in the adult.  
 The nest was made of sticks and mud, and was  
 - 12/2/83: The first egg - and the only one -  
 was found in the nest. - 15/6 In 2 eggs  
 from the nest. - 27/6/83 and the nest was found  
 in the nest. The nest was found in the nest.  
 The nest was found in the nest.

*Lanius borealis*

*Samolus purpureus*  
7/5 88 got one from ~~John~~ ~~Gardner~~, ~~Saranac~~ ~~Co.~~

*Chalix pauciflora*  
26/4 88 - 3 lb. at ~~Saranac~~, ~~Co.~~ 9.  
2/5 88 Early in the morning plenty of it from ~~Co.~~  
- Cooks thinks that if *P. p.* occurs at  
all in ~~Co.~~, it will only be very  
rare, he has seen it in two places  
and never saw it before. It was found  
there in a bush in ~~Co.~~. The ~~Co.~~  
say that it is not abundant in ~~Co.~~

Arthur

*Spilothamnops pictus*.

7589. Swampy bank of river seen at the  
today, Oct. -

*Cyprus Chambe*

Mr. W. H. Jones, that he has seen one  
at the 22 to 24 1738 - 14383. several  
at the 22 - several hundred individuals  
at Topkay 4/683 in smaller or larger flocks.

*Spilothamnops pictus*



Uria arctica


No birds were seen on the 1st of  
Aug. 1880. Spent 1st & 2nd Sept. at  
Long Island, Alaska, where we were to stay.  
— When sitting on the rock shelves, the  
Arctic <sup>usually</sup> turn their breast to the rock wall and then  
back to the sea, while Lunda cirrhata and  
Fratercula corniculata front the sea. ~~usually~~ Their  
position of the latter is surely upright although  
not so straight as that of the Uria —  
"true Lunda".

*Metacilla leucophaea*

*Chrysomelidae*

*Thylophora brenckei*

1881



THOMAS GROOM & CO.  
STATIONERS,  
(82 State Street)  
BOSTON.



